

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.



EPA Reg. No. 70051-86

ACTIVE INGREDIENT: *Bacillus thuringiensis* subspecies *kurstaki* strain EG7841 solids, spores and Lepidopteran active toxins*: 40.0%

OTHER INGREDIENTS: 60.0%
TOTAL: 100.0%

*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

PESTS

For control of light brown apple moth (LBAM) (*Epiphyas postvittana*), banana moth (*Opogona sacchari*), European grapevine moth (*Lobesia botrana*) and European pepper moth (*Duponchelia fovealis*).

For light brown apple moth: Apply when newly hatched larvae appear and before leaves are rolled or webbing is significant.

For banana moth: Drench bark to newly emergent shoots following pruning or apply to susceptible plant tissues when banana moth larvae are active.

For European grapevine moth: Apply at blackhead egg stage or when larvae are newly hatched before leaves are rolled, or larvae have entered fruit.

For European pepper moth: Begin applications at egg lay and continue at 3-5-day intervals throughout larval feeding period.

For all pests: Continue applying as part of a normal spray program until pest is adequately controlled. Apply when caterpillars are actively

feeding. To be effective, Crymax spray must be deposited at the larval feeding site. Crymax can be applied by ground or air in water sufficient to insure thorough and even coverage. Thorough and uniform crop coverage is required for adequate insect control. Applications at higher water volumes have demonstrated improved control of targeted pests. Early morning or evening applications, when air is calm, are generally best for aerial applications.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **For Utah only, this recommendation is not required to be in the possession of the user at the time of pesticide application. For all states other than Utah, this recommendation must be in the possession of the user at the time of pesticide application.**

Continued on reverse side

Crymax[®]

BIOINSECTICIDE

APPLICATION RATE

Apply Crymax at 0.5 to 2.0 pounds per acre to the following crops:

Vegetable Crops:

Artichokes, arugula, asparagus, beans, beets, bok choy, broccoli, brussel sprouts, cabbage, cardoni, carrots, cauliflower, celeriac, celery, chick peas, chicory, Chinese cabbage, collards, cucumber, cucurbits, dry bulb onions, eggplant, escarole, endive, garlic, green onions, greens (beets, China, dandelion, mustard, turnip), horseradish, kale, kohlrabi, leeks, lentils, lettuce (head, leaf, romaine), malanga, melons (cantaloupe, crenshaw, honeydew, muskmelon, watermelon, etc.), napa, okra, onions, parsley, parsnips, peas, peppers, potatoes, pumpkins, radishes, rutabaga, salsify, shallots, soybean foliage, spinach, squash, sugarbeets, sweet potatoes, swiss chard, tomatoes, turnips

Herbs and Spices:

Basil, chives, cilantro, dill, oregano, peppermint, thyme

Fruit, Nut and Vine Crops:

Pome and stone fruits: apples, apricots, cherries, nectarines, peaches, pears, plums, prunes, quince

Nut trees: almonds, chestnuts, filberts, pecans, pistachios, walnuts

Citrus

Small fruit and berries: blackberries, blueberries, boysenberries, cranberries, currants, loganberries, raspberries, strawberries

Grapes

Tropical and other fruit: avocados, bananas, kiwi, persimmons, pomegranate, pineapple, tropical fruits

Field Crops:

Canola/rape seed, evening primrose, meadow foam, corn (field, sweet popcorn, seed), cotton, hops, jojoba, peanuts, rice, safflower, small grains (barley, oats, rye, wheat, etc.), sorghum, soybeans, sunflowers, tobacco

Commercial Flowers and Ornamental Plants:

Bedding plants, flowers (greenhouse and field), greenhouse ornamentals, greenhouse vegetables, container stock

Forest, Shade Tree and Nursery Stock:

Forest trees, shade trees, nursery trees