

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.

Cueva®

FLOWABLE LIQUID COPPER FUNGICIDE

EPA Reg. No. 67702-21

ACTIVE INGREDIENT: Copper Octanoate (Copper Soap): 10.0%

CAS Reg. No. 20543-04-8

OTHER INGREDIENTS: 90.0%

TOTAL: 100.0%

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROPS

Vegetables, herbs, field crops, nuts, fruits and berries

PESTS

For control of citrus canker, *Xanthomonas citri* subsp. *citri*.

RATE

Mix 0.5 to 2 gallons of Cueva fungicide concentrate with 100 gallons of water. Apply 50 to 100 gallons of diluted spray per acre.

APPLICATION

For application by aircraft, apply 5 to 25 gallons of diluted spray to one acre. For best control, begin treatment 2 weeks before disease normally appears or when weather forecasts predict a long period of wet weather. Alternatively, begin treatment when disease first appears and repeat at 7- to 10-day intervals for as long as needed. Use the 2-gallon rate of Cueva fungicide concentrate, applied every 7 days or less, following heavy rain or when the amount of disease is increasing rapidly. If possible, time applications so that 12 hours of dry weather follow application.

Citrus Crops

For citrus canker apply every 3 weeks after petal fall until harvest.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **For Utah only, this recommendation is not required to be in the possession of the user at the time of pesticide application. For all states other than Utah, this recommendation must be in the possession of the user at the time of pesticide application.**