

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.

Double Nickel® LC

BIOFUNGICIDE

EPA Reg. No. 70051-107

Aqueous Suspension Biofungicide/Bactericide

ACTIVE INGREDIENT: *Bacillus amyloliquefaciens* strain D747*: 98.85%

OTHER INGREDIENTS: 1.15%

TOTAL: 100.0%

*Contains a minimum of 1×10^{10} colony-forming units (cfu) per milliliter.

FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY

Expires 12/31/2020

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

PEST or DISEASE

For suppression of Root Knot (*Meloidogyne* spp.), Cyst (*Globodera* spp.), Lesion (*Pratylenchus* spp.), Ring (*Mesocriconema* spp.) and Spiral (*Helicotylenchus* spp.) nematodes.

Nematode suppression has been seen when Double Nickel LC is applied as part of a soil disease management program.

SOIL APPLICATION

For control of soilborne diseases infecting seeds, seedlings, roots, crown, stems, or other plant parts below ground or in contact with soil: Apply Double Nickel LC at 0.5 to 4.5 pints per acre. Mix the required amount in sufficient water to apply by one of the following methods:

- Soil drench applied to transplants in flats or pots in the greenhouse or nursery any time prior to transplanting (refer to label for instructions).
- Soil drench at transplanting, using a "water wheel" injector, spray nozzles/hoses, or other method to drench each root ball and/or planting hole.

- Soil or seed line drench, or banded spray (in-furrow) at planting (refer to label for specific instructions).

FOLLOW UP

Follow-up (post-planting) preventive applications can be made every 2-4 weeks by one or more of the following methods, if needed:

- Drip (trickle) or any type of sprinkler irrigation, any time after planting or transplanting. See Chemigation Instructions for additional information.
- Spray directly onto the soil surface and/or lower plant parts. If targeting root disease, follow immediately with sufficient overhead sprinkler irrigation to move Double Nickel LC to the root zone.
- Injection directly into the rooting zone using shanks or similar equipment.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**