

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.

MeloCon® WG

BIOLOGICAL NEMATOCIDE

EPA Reg. No. 72444-2

ACTIVE INGREDIENT: *Paecilomyces lilacinus* strain 251*: 6.0%

OTHER INGREDIENTS: 94.0%

TOTAL: 100.0%

*Contains a minimum of 1 X 10¹⁰ spores per gram of product

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROPS

Vegetables, citrus, peaches, bananas, grapes, tree nuts, strawberries, ornamentals, pineapples, and tobacco

PESTS

Plant parasitic nematodes, including:

- Awl nematode (*Dolichodorus heterocephalus*)
- Burrowing nematode (*Radopholus similis*)
- Citrus nematode (*Tylenchulus semipenetrans*)
- Cyst nematode (*Heterodera* and *Globodera*)
- False root knot nematode (*Nacobus*)
- Lance nematode (*Hoplolaimus columbus*)
- Lesion nematode (*Pratylenchus*)
- Needle nematode (*Longidorus* and *Paralongidorus*)
- Reniform nematode (*Rotylenchulus reniformis*)
- Ring nematode (*Criconemoides*, *Criconebella* and *Mesocriconebella*)
- Root knot nematode (*Meloidogyne*)
- Spiral nematode (*Helicotylenchus*)
- Stem nematode (*Ditylenchus dipsaci*)
- Sting nematode (*Belonolaimus longicaudatus*)
- Stubby root nematode (*Trichodorus* and *Paratrachodorus*)
- Stunt nematode (*Tylenchorhynchus*)

RATES

2 to 4 pounds per acre

USE RECOMMENDATIONS

For field treatment:

Mix required amount of MeloCon WG in water and apply to soil by drenching, shank injection, or through drip or sprinkler irrigation. Apply in sufficient water to move the product into the root zone.

Apply to nematode-infested soil 14 days before planting or transplanting. Apply to recently fumigated soil at planting or transplanting.

Repeat applications at 6 week intervals as needed to suppress plant parasitic nematode populations during the crop period.

Application efficiency may be enhanced by inclusion of a soil wetting agent to help the spores penetrate soil to the root zone.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **For Utah only, this recommendation is not required to be in the possession of the user at the time of pesticide application. For all states other than Utah, this recommendation must be in the possession of the user at the time of pesticide application.**

MeloCon is a registered trademark of Bayer CropScience Biologics GmbH.