

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.

OSO™ 5% SC
FUNGICIDE

EPA Reg. No. 68173-4-70051

ACTIVE INGREDIENT: Polyoxin D zinc salt: 5.0%

OTHER INGREDIENTS: 95.0%

TOTAL: 100.0%

Contains 7.03 ounces of active ingredient per gallon.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROPS

Pome Fruits including apple, crabapple, loquat, mayhaw, pear, and quince.

PESTS or DISEASE

For control of Sooty blotch, Flyspeck, Bitter rot, Alternaria rot and suppression of Cedar apple rust and White rot.

RATE and APPLICATION

Begin applications of OSO 5% SC Fungicide at 6.5 oz/acre prior to disease development and continue on a 7 to 10 day interval. OSO 5% SC Fungicide may be applied anytime from green-tip up to day of harvest.

States

CT, DE, MA, MD, ME, MI, NC, NH, NJ, NY, OH, PA, VA, VT, WV.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**