

OSO™ 5%SC FUNGICIDE

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

OSO™ 5% SC Fungicide

EPA Reg. No. 68173-4-70051

ACTIVE INGREDIENT:

Polyoxin D zinc salt5.0%

OTHER INGREDIENTS:95.0%

TOTAL:100.0%

Contains 7.03 ounces of active ingredient per gallon.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP: Pome Fruits (including apple, crabapple, loquat, mayhaw, pear, and quince).

PEST or DISEASE:

For control of Sooty blotch, Flyspeck, Bitter rot, Alternaria rot and suppression of Cedar apple rust & White rot.

APPLICATION INSTRUCTIONS:

Begin applications of OSO 5% SC Fungicide at 6.5 oz/acre prior to disease development and continue on a 7 to 10 day interval. OSO 5% SC Fungicide may be applied anytime from green-tip up to day of harvest.

STATES: CT, DE, MA, MD, ME, MI, NC, NH, NJ, NY, OH, PA, VA, VT, WV

All applicable directions, application rates, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels.

This recommendation must be in the possession of the user at the time of pesticide application.