

**FIFRA Section 2(ee)
Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

SoilGard[®]12G
MICROBIAL FUNGICIDE

EPA Reg. No. 70051-3

PESTS For control of *Sclerotinia* sp., *Phytophthora capsici*, *Sclerotium* and *Fusarium* root and crown rots.

DIRECTIONS FOR USE

CROPS	RATE	APPLICATION
Transplanted crops such as fruiting and leafy vegetables, melons, strawberries, field grown ornamental plants, and nonbearing fruit/nut trees and vines.	Before Transplanting 1.5 to 2 lbs per 100 gallons of water	Apply 4 fl. oz. of finished drench per plant in flats (unit trays) or pots within 1 week before transplanting into the field (during the hardening-off phase). For densely seeded trays or flats, apply 100 gallons of finished drench per 800 square feet. SoilGard 12G can also be applied in furrow immediately before transplanting as described for seed furrows in the next section.
	At Transplanting 1.5 to 2 lbs per 100 gallons of water	Apply 4-8 fl. oz. of finished drench (depending on the size of the root ball) in each transplant hole using a water wheel/shank injector, watering can or low pressure sprayer with flood/drench nozzles.
	After Transplanting 2 to 10 lbs per acre	Repeat (maintenance) applications can be made as needed through directed spray or drench at the base of each plant, or through irrigation system. (See "For Application Through Irrigation Systems" on the current SoilGard 12G label).
Seed furrows or planting beds such as potatoes and other tubers, bulbs, and direct-seeded crops.	2 to 10 lbs per acre	Apply in 50 to 100 gallons of water per acre as banded drench in-furrow at or immediately before planting, and before covering seed or seed pieces with soil. Apply a band of sufficient width to cover the seed furrow and initial rooting zone (usually 4-8 inches, centered on the furrow). Drench can also be applied as a seed line treatment using shank or other injection equipment in sufficient water to treat both the seed furrow and initial rooting zone.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. This recommendation must be in the possession of the user at the time of pesticide application.